



## Environmental impacts of precision feeding programs applied in Brazilian pig production

POMAR, Candido<sup>1</sup>; ANDRETTA, Ines<sup>2</sup>; HAUSCHILD, Luciano<sup>3</sup>; KIPPER, Marcos<sup>2</sup>; PIRES, Paula Gabriela<sup>2</sup>

<sup>1</sup>Agriculture and Agri-Food Canada, Sherbrooke, Quebec, Canada; Candido.Pomar@agr.gc.ca.

<sup>2</sup>Universidade Federal do Rio Grande do Sul, Porto Alegre, Rio Grande do Sul, Brazil.

<sup>3</sup>Universidade Estadual Paulista, Jaboticabal, São Paulo, Brazil.

A paper from the Proceedings of the  
14<sup>th</sup> International Conference on Precision Agriculture  
June 24 – June 27, 2018  
Montreal, Quebec, Canada

**Abstract.** *This study was undertaken to evaluate the effect that switching from conventional to precision feeding systems during the growing-finishing phase would have on the potential environmental impact of Brazilian pig production. Standard life-cycle assessment procedures were used, with a cradle-to-farm gate boundary. The inputs and outputs of each interface of the life cycle were organized in a model. Grain production was independently characterized in the Central-West and South regions of Brazil, while the pigs were raised in the South region. Three feeding programs were applied for growing-finishing pigs: CON, conventional phase feeding by group; PFG, precision daily feeding by group (whole herd fed the same daily adjusted diet); and PFI, precision daily feeding by individual (diets adjusted daily to match individual nutrient requirements). Raising pigs (1 t pig body weight at farm gate) in South Brazil under the CON feeding program using grain cultivated in the same region led to emissions of 1,840 kg of CO<sub>2</sub>-eq, 13.1 kg of PO<sub>4</sub>-eq, and 32.2 kg of SO<sub>2</sub>-eq. Simulations using grain from the Central-West region showed a greater climate change impact. Compared with the previous scenario, a 17% increase in climate change impact was found when simulating with soybeans produced in Central-West Brazil, while a 28% increase was observed when simulating with corn and soybeans from Central-West Brazil. Compared with the CON feeding program, the PFG and PFI programs reduced the potential environmental impact. Applying the PFG program mitigated the potential climate change impact and eutrophication by up to 4%, and acidification impact by up to 3% compared with the CON program. Making a further adjustment by feeding pigs according to their individual nutrient requirements mitigated the potential climate change impact by up to 6% and the potential eutrophication and acidification impact by up to 5% compared with the CON program. The greatest environmental gains associated with the adoption of precision feeding were observed when the diet combined soybeans from Central-West Brazil with corn produced in Southern Brazil. The results of this study clearly show that precision feeding is an effective approach for improving the environmental sustainability of Brazilian pig farming.*

**Keywords.** *life-cycle assessment, nutrient requirements, precision farming, swine.*

Commented [IA1]: Limit: 350 words long

---

The authors are solely responsible for the content of this paper, which is not a refereed publication. Citation of this work should state that it is from the Proceedings of the 14th International Conference on Precision Agriculture. EXAMPLE: Lastname, A. B. & Coauthor, C. D. (2018). Title of paper. In Proceedings of the 14th International Conference on Precision Agriculture (unpaginated, online). Monticello, IL: International Society of Precision Agriculture.

---

## Introduction

Many studies have found that nutrient efficiency can be improved by adjusting the nutrient supply to more closely match the animals' individual requirements (Ferket *et al.*, 2002; Pomar *et al.*, 2014). Nutritional requirements change dynamically over time and also vary among animals, even in age- or sex-homogeneous populations (Pomar *et al.*, 2003; Brossard *et al.*, 2009). However, current group phase-feeding programs do not account for variations among individuals, and feeds are usually formulated to optimize the performance of the population, which means that most pigs receive more nutrients than they actually need (Hauschild *et al.*, 2010).

Although dealing with variability in nutritional requirements is a difficult task, precision farming techniques may provide a solution by allowing pigs to be managed and fed individually using diets adjusted in real-time to match their nutritional requirements (Pomar *et al.*, 2009; Hauschild *et al.*, 2012). This approach represents a paradigm shift in pig feeding, since the optimal dietary nutrient level is no longer considered a static population attribute, but rather a dynamic process that evolves independently for each animal (Pomar *et al.*, 2014).

Precision feeding techniques reduced lysine intake and nitrogen excretion without compromising pig performance in previous studies (Andretta *et al.*, 2014; Andretta *et al.*, 2016). Despite the potential benefits of precision feeding (Wathes *et al.*, 2008), the global impact of these practices on sustainability merits further investigation.

Feeds contribute highly to the environmental impacts of livestock products. This had already been stated for Brazilian pig production in research developed using life-cycle assessment (Spies, 2009; Cherubini, *et al.*, 2015a, 2015b). Previous studies also reported that improving feeding practices may mitigate the environmental footprint of producing pigs in Brazil (Kebreab, *et al.*, 2016; Monteiro *et al.*, 2016). The present study was therefore undertaken to evaluate the effect that switching from a conventional feeding system to precision feeding programs during the growing-finishing phase would have on the potential environmental impact of pig production.

## Material and methods

Environmental impacts were evaluated according to the life-cycle assessment (LCA) standards using the approach described by Guinée (2002) based on four interrelated steps: goal and scope definition; life cycle inventory; life cycle impact assessment, and interpretation.

Brazil was chosen for this simulation because it is a large producer and exporter of pork. Moreover, despite the importance of the pig sector in developing countries, most studies on the environmental cost of pork have been developed based on European conditions, with limited applicability to other major pig production regions.

The major stages considered in the model were the production of feed ingredients from plant sources (corn and soybean meal), the production of other feed ingredients (amino acids, limestone, dicalcium phosphate, salt, and vitamin-mineral premix), drying and processing in the feed industry, transportation, and animal rearing (Figure 1). The functional unit used to study the environmental cost of grain production was 1 t of each grain at the feed factory. The functional unit used to study potential environmental impacts related to the feeds was 1 t of rations produced and delivered to the farm. Lastly, when assessing the cradle-to-farm gate model, environmental costs were simulated using 1 t body weight of finished pig (120 kg of individual slaughter weight) at farm gate as the functional unit.




Fig 1. Basic flowchart of the production system.

### Description of the pig production system

This case study was developed for a complete farrow-to-finish unit with a confinement operation (buildings with concrete floors) located in the municipality of Concórdia, in the state of Santa Catarina, Brazil. This scenario was chosen because it is an important traditional pig production region in South Brazil. The transport route between the farm and the feed factory was defined as the average distance between these productive units in Brazil, namely 35 km (Talamini *et al.*, 2006).

### Description of the production system for grains and other feed ingredients

Grain production was independently characterized in the Central-West region (CW) and South region (SO) of Brazil. The crop farm locations were chosen based on the ranking of the major corn- and soybean-producing municipalities in each region (IBGE, 2014). Agricultural practices for grain production and models used to calculate the emissions in the crop systems were adapted from Alvarenga (2010); Alvarenga *et al.* (2012) and Prudêncio da Silva *et al.* (2010). Land transformation was estimated based on the data provided by Alvarenga (2010) and the methodology described by Prudêncio da Silva *et al.* (2010). Grain yield data were obtained for each city from the Brazilian Institute of Geography and Statistics (IBGE, 2014).

The environmental cost of grain production varies greatly among the Brazilian regions. Consequently, three geographic scenarios were simulated based on different crop cultivation locations: CW-CW, in which only grain grown in CW were used to produce the feed; CW-SO, in which soybeans from CW and corn from SO were used to produce the feed; and SO-SO, in which only grain from SO were used to produce the feed. The scenarios differed mainly in terms of road transportation distances, agricultural practices, and deforestation impact on newly opened agricultural frontiers (deforestation was assumed for the CW scenario only because it occurred many decades ago in the SO region).

The scope of synthetic amino acid production was adapted from Mosnier *et al.* (2011), distinguishing amino acids produced by chemical synthesis (DL-methionine) from those produced by fermentation (L-lysine and L-threonine). Other feed ingredients were based on the Ecoinvent database (v. 3.0, Swiss Centre for Life Cycle Inventories, Dübendorf, Switzerland). The environmental impact of vitamin-mineral trace elements was assumed to be equal to that of limestone.

Grain processing and storage conditions were adapted from previous reports (van Zeist *et al.*, 2012a and 2012b). Grain (from the farm to the feed factory), other ingredients (from industry to the feed factory) and feeds (from the feed factory to the pig farm) were assumed to have been transported by truck. Google Earth software (Google Inc., Mountain View, CA) was used to estimate road distances. Information from the Agri-footprint database (v. 1, Blonk Agri-footprint BV, Gouda, The Netherlands) was used to simulate the impact of transportation.

### Feeding practices and animal performance

The characterization of animal performance and feeding practices was mainly focused on the growing-finishing pigs, since the adoption of conventional and precision feeding programs has previously been tested in this rearing phase. Other production stages (breeding-gestation, farrowing, and nursery phases) were included in the scope of this study, but similar production conditions in terms of management and feeding practices were considered for all the scenarios being compared. Culled sows were considered a co-product of piglet production.

Ingredients commonly used in Brazilian industrial pig farming were used to formulate all the feeds. The formulas used soybean meal as the major protein source and corn as the major energy supply. Wheat was used only in the gestation feed. Simple formulas were simulated in this study because the use of alternative ingredients vary greatly among scenarios (e.g. among industries, geographical regions, seasons or years) according to their availability and cost. The least-cost formulation method was applied based on annual average prices (CEPEA-ESALQ, 2014). The nutritional composition of the ingredients (net energy, standardized ileal digestible amino acids, digestible phosphorus, and total calcium) was determined using EvaPig software (v. 1.3.1.4, INRA, Saint-Gilles, France).

Feeds for breeding animals and for nursery piglets were formulated based on the nutritional requirements and the feeding program proposed in the Brazilian Tables for Poultry and Swine (Rostagno *et al.*, 2011). Average performance indicators for describing the breeding-gestation, farrowing, and nursery phases were obtained from industry reports (Agriness, 2013).

Data on a reference population of growing-finishing pigs (130 animals of a high-performance genotype previously described by Andretta *et al.* (2014) and Andretta *et al.* (2016)) were used to compare the feeding programs in terms of performance and nutrient excretion. Average performance data were applied to the conventional and precision feeding programs, since these treatments did not influence feed intake or body weight gain in the previous studies. However, individual nitrogen and phosphorus retention data were considered when estimating nutrient excretion in each feeding program.

Individual data collected on the reference population was also used to estimate the nutritional requirements used to define all the feeding programs. This procedure was based on the required dietary concentration of lysine, which was estimated for each pig using a previously described mathematical model (Hauschild *et al.*, 2012) based on individual daily feed intake and weekly body weight information. In the model, the empirical component estimated the expected body weight, feed intake, and daily gain for the next day, and the mechanistic component then used these three estimates to calculate, according to a factorial method, the optimal concentration of amino acids that should be offered that day to each pig in the herd so as to meet their requirements. Based on the estimated individual nutrient requirements, three feeding programs for growing-finishing pigs were proposed and simulated in the current study: conventional phase feeding (CON), precision daily feeding by group (PFG), and precision daily feeding by individual (PFI).

The CON treatment consisted of a three-phase feeding program in which the whole herd received the same feed during each 28-d phase. The complete feeds used in this program were formulated to satisfy the requirements of the 80th-percentile pig in the population on the first day of each phase, as suggested by Hauschild *et al.* (2010) to maximize population responses for weight gain.

Two blends of premixes (named A and B) were adjusted to match the lysine requirements of the pigs assigned to the daily phase-feeding programs. These premixes differed in their nutrient concentrations: premix A was formulated with a high nutrient density (to satisfy the estimated requirements of the most demanding pig at the beginning of the growing period), and premix B formulated with a low nutrient density (to satisfy the estimated requirements of the least demanding pig at the end of the finishing period).

The optimal dietary nutrient concentration in the PFG program was calculated for each day based on an animal whose requirement placed it in the 80th percentile of the population (Hauschild *et al.*, 2010). Lastly, the optimal dietary lysine concentration in the PFI program was calculated for each day based on individual nutrient requirements (Hauschild *et al.*, 2012). In this program, each pig would be fed with a blend of premixes A and B that was adjusted on a daily basis to match its individual requirements.

Manure management was considered and includes liquid manure storage in slurry tanks, transport, and field application for fertilization in replacement of fertilizers (Alvarenga, 2010). When possible, calculations were developed considering that feeding practices affected manure composition, which was calculated based on the nutrient input balance in the feed and the nutrient retention rate measured in the reference population (Andretta *et al.*, 2014; Andretta *et al.*, 2016). Emissions of NH<sub>3</sub> related to manure management were estimated according to Eriksson *et al.* (2005), while the Intergovernmental Panel on Climate Change methodology (IPCC, 2006) was used to estimate the emissions of N<sub>2</sub>O (direct and indirect; Tier 2) and CH<sub>4</sub> (from enteric fermentation and waste management; Tier 1).

### **Modeling environmental impacts**

Inputs and outputs were defined for each step of the life cycle and organized in a model using SimaPro software (v. 8.0.3.14, PRE Consultants, Amersfoort, The Netherlands). Environmental impacts related to medication and capital assets (machinery, equipment, and buildings) were not considered in the model. Slaughtering procedures were also excluded, because a previous study found that feeding programs had no effect on carcass or meat traits (Andretta *et al.*, 2014). The allocation of environmental burdens to by-products (e.g. soybean meal and oil) was based on economic criteria.

Climate change, eutrophication, and acidification were the chosen environmental impact categories. Results were obtained for each environmental impact category, stating the resources used in each production system and the aggregate emissions of each substance with the characterization factor in the impact categories. The CML-IA baseline method was used for the calculations.

### **Assessment of the environmental impact of feeds adjusted to individual nutrient requirements**

The impact of heterogeneous nutritional requirements on the environmental cost of feeds was analyzed using the reference population of growing-finishing pigs. In this sensitivity analysis, the entire reference population was used to generate a database with estimated nutritional requirements for each pig per day. Feeds were adjusted to meet the daily and individual nutrient requirements, generating one dietary formula per pig for each day. Lastly, the environmental cost of each adjusted feed was estimated and analyzed in terms of population heterogeneity. Regression equations were also used to analyze the effect of dietary nutrient composition on the potential environmental impact associated with the feeds.

## Results and discussion

### Assessment of the environmental impact of grain production

Life-cycle assessment of grain production is an essential part of evaluating the environmental impact of pig production. Crop management practices and expansion rates vary greatly among Brazilian regions. The production of soybean meal (functional unit: 1 t at the feed factory) in SO was associated with the emission of 533 kg of CO<sub>2</sub>-eq, 5.82 kg of PO<sub>4</sub>-eq, and 2.62 kg of SO<sub>2</sub>-eq. In comparison with the SO scenario, soybean meal from CW showed higher climate change (+108%) and acidification (+107%) impacts, while producing a lower eutrophication impact (-3%). Production of corn (1 t at feed factory) in the SO scenario led to the emission of 491 kg of CO<sub>2</sub>-eq, 3.78 kg of PO<sub>4</sub>-eq, and 9.98 kg of SO<sub>2</sub>-eq. In the CW scenario, corn showed a higher impact with regard to climate change (+22%) and eutrophication (+3%) but a lower impact with regard to acidification (-11%) compared with the SO scenario.

Variations between the Brazilian CW and SO scenarios have already been reported in relation to the environmental impact associated with grain (Prudêncio da Silva *et al.*, 2010; Mosnier *et al.*, 2011) and feed production (Alvarenga *et al.*, 2012). These differences are related to specific crop cultivation practices applied in each region under study. Longer transportation distances also contributed to the higher environmental impact of grain from CW. In addition, grain cultivated in CW was associated with the recent conversion (within the past 30 years) of natural forests into cropland, which was not observed in the SO scenario.

### Assessment of the environmental impact of Brazilian pig diets

The potential environmental impacts of conventional diets for growing-finishing pigs are presented in Table 1. Similar environmental impacts were reported by Alvarenga *et al.* (2012) for the diets of broiler chickens produced in Brazil. Cherubini *et al.*, 2015b also reported comparable impacts in terms of the carbon footprint when assessing diets for finishing pigs in Brazil. The climate change and eutrophication impacts estimated in the current study were also comparable to those obtained by van der Werf *et al.* (2005) and Mosnier *et al.* (2011), who assessed diets for finishing pigs produced in France using Brazilian soybeans.

**Table 1. Performance, feeding costs and nitrogen balance of pigs fed in 3-phases (3P), commercial (COM), multi-phases per group (MPG) and individual multi-phases (MPI) feeding programs (trial 1).**

Response	Premix <sup>1</sup>		Conventional feed <sup>2</sup>		
	A	B	Phase I	Phase II	Phase III
<b>SO-SO scenario<sup>3</sup></b>					
Climate change, kg CO <sub>2</sub> -eq	557	520	548	543	537
Eutrophication, kg PO <sub>4</sub> -eq	4.35	3.79	4.31	4.22	4.14
Acidification, kg SO <sub>2</sub> -eq	8.21	9.91	9.46	9.79	10.1
<b>CW-SO scenario<sup>4</sup></b>					
Climate change, kg CO <sub>2</sub> -eq	711	532	691	664	637
Eutrophication, kg PO <sub>4</sub> -eq	4.30	3.79	4.26	4.18	4.11
Acidification, kg SO <sub>2</sub> -eq	8.97	9.96	9.02	9.18	9.34
<b>CW-CW scenario<sup>5</sup></b>					
Climate change, kg CO <sub>2</sub> -eq	786	638	769	746	724
Eutrophication, kg PO <sub>4</sub> -eq	4.37	3.88	4.33	4.25	4.18
Acidification, kg SO <sub>2</sub> -eq	8.24	8.94	8.27	8.38	8.49

<sup>1</sup> Premix: Used in the precision feeding programs and formulated to contain a high (Premix A) or low (Premix B) density of crude protein, amino acids, and phosphorus.

<sup>2</sup> Complete feed: Used in the conventional feeding program.

<sup>3</sup> SO-SO scenario: Soybeans and corn produced in South Brazil.

<sup>4</sup> CW-SO scenario: Soybeans produced in Central-West Brazil and corn produced in South Brazil.

<sup>5</sup> CW-CW scenario: Soybeans and corn produced in Central-West Brazil.

Due to their inclusion in the feed formula, corn and soybean meal were the ingredients with the highest contribution to the environmental impact. On average, corn production accounted for 72% of the total potential climate change impact attributed to commercial feeds in the SO-SO scenario, while 22% of the impact was due to the soybean chain, 3% to dicalcium phosphate, and

*Proceedings of the 14<sup>th</sup> International Conference on Precision Agriculture  
June 24 – June 2, 2018, Montreal, Quebec, Canada*

2% to synthetic amino acid production. Very similar patterns were observed for other environmental impact categories and geographical scenarios.

Feeds formulated for growing pigs (Phase 1) showed a higher impact in terms of climate change (SO-SO: +2%, CW-SO: +8%, CW-CW: +6%) and eutrophication (SO-SO: +4%; CW-SO: +4%, CW-CW: +3%) compared with feeds for finishing animals (Phase 3). Similar patterns were observed among the premixes. A higher impact in terms of climate change (SO-SO: +7%, CW-SO: +25%, CW-CW: +19%) and eutrophication (SO-SO: +13%, CW-SO: +12%, CW-CW: +11%) was estimated for premix A (with a high density of crude protein and other nutrients) than for premix B (with a low density of protein and other nutrients).

The nutritional requirements used to formulate a feed may influence its environmental impact. To test this hypothesis, a simulation was performed involving feeds adjusted to individual nutrient requirements that had been estimated using the reference population of growing-finishing pigs. The variability in terms of climate change impact among simulated feeds is due to the changes in the incorporation of high-impact ingredients, which is a response to the nutrient requirement variability among pigs. However, this impact varied depending on the geographical scenario. The intraday average range in the potential climate change impact associated with the production of 1t of feeds was 36.3 kg of CO<sub>2</sub>-eq in the SO-SO scenario, while it was estimated at 145.2 kg of CO<sub>2</sub>-eq in CW-CW and 175.6 kg of CO<sub>2</sub>-eq in CW-SO. In the same comparison, the estimated intraday interquartile range (25th to 75th percentile) was 8.1 kg of CO<sub>2</sub>-eq in the SO-SO scenario, while it was 32.3 kg of CO<sub>2</sub>-eq in CW-CW and 39.0 kg of CO<sub>2</sub>-eq in CW-SO. The intraday coefficient of variation for the potential climate change impact of population-adjusted feeds was also lower in the SO-SO scenario (1.4% on average) than in the CW-CW (4.2%) and CW-SO (5.9%) scenarios.

The simulation also highlighted a reduction over time in the potential climate change impact associated with pig feed production that was closely related to the reduction in the expected dietary nutrient levels. In the simulated population, reducing the dietary standardized ileal digestible lysine level by one percentage point led to a reduction of up to 194.7 kg of CO<sub>2</sub>-eq per t of feed in the CW-SO scenario. Several studies have reported that it may be possible to mitigate the environmental impact associated with pig feeds by reducing the dietary crude protein level and, consequently, the use of soybean products in the feed formulas (Eriksson *et al.*, 2005; Mosnier *et al.*, 2011; Meul *et al.*, 2012; Cherubini, *et al.*, 2015b). In most scenarios, as in CW, soybean production has a higher environmental cost than other feed ingredients, especially when compared with domestic alternatives (Eriksson *et al.*, 2005; Monteiro *et al.*, 2016). Applying precision feeding techniques is a way to reduce the protein content in the diets and consequently the use of soybean products in pig feeding. This condition support the hypothesis that precision feeding techniques may be an effective alternative for reducing the environmental impact of pig production, as previously indicated by Monteiro *et al.* (2016). In addition, variations among the intercepts found for the studied regions may also indicate that the environmental impact of changing feeding programs depends on the production scenario used for the simulation.

### **Assessment of the environmental impact of conventional Brazilian pig production**

The potential environmental impact of finished pigs is presented in Table 2. Raising pigs (1 t body weight pig at farm gate) in South Brazil according to the CON feeding program and using grain cultivated in the same region (SO-SO scenario) led to the emission of 1,840 kg of CO<sub>2</sub>-eq, 13.1 kg of PO<sub>4</sub>-eq, and 32.2 kg of SO<sub>2</sub>-eq.

The environmental costs of pig production in the state of Santa Catarina were previously simulated by Spies (2009) as 1,720 kg of CO<sub>2</sub>-eq, 9.55 kg of PO<sub>4</sub>-eq, and 19.8 kg of SO<sub>2</sub>-eq per ton of finished pig (body weight at slaughter). Cherubini *et al.* (2015a) also studied the environmental profile of swine production in South Brazil and estimated emissions at 3,500 kg of CO<sub>2</sub>-eq per ton of swine carcasses produced. The variation among results is partially explained by the difference in functional units and boundaries used in the latter study (slaughtering



procedures were accounted for and a carcass yield of 73.9% was applied).

**Table 2. Performance, feeding costs and nitrogen balance of pigs fed in 3-phases (3P), commercial (COM), multi-phases per group (MPG) and individual multi-phases (MPI) feeding programs (trial 1).**

Response	Feeding Program		
	Conventional Phase Feeding by Group	Precision Daily Feeding	
		By Group	By Individual
<b>SO-SO scenario<sup>1</sup></b>			
Climate change, kg CO <sub>2</sub> -eq	1,840	1,811	1,783
Eutrophication, kg PO <sub>4</sub> -eq	13.1	12.7	12.4
Acidification, kg SO <sub>2</sub> -eq	32.2	31.4	31.0
<b>CW-SO scenario<sup>2</sup></b>			
Climate change, kg CO <sub>2</sub> -eq	2,160	2,079	2,030
Eutrophication, kg PO <sub>4</sub> -eq	13.0	12.6	12.3
Acidification, kg SO <sub>2</sub> -eq	33.8	32.8	32.2
<b>CW-CW scenario<sup>3</sup></b>			
Climate change, kg CO <sub>2</sub> -eq	2,361	2,300	2,252
Eutrophication, kg PO <sub>4</sub> -eq	13.2	12.7	12.5
Acidification, kg SO <sub>2</sub> -eq	31.8	30.8	30.1

<sup>1</sup> SO-SO scenario: Soybeans and corn produced in South Brazil.

<sup>2</sup> CW-SO scenario: Soybeans produced in Central-West Brazil and corn produced in South Brazil.

<sup>3</sup> CW-CW scenario: Soybeans and corn produced in Central-West Brazil.

The geographical scenarios simulated for grain production had a minor effect on emissions of PO<sub>4</sub>-eq and SO<sub>2</sub>-eq. The highest variations between the scenarios for eutrophication (-2%) and acidification (+6%) were found between the CW-SO and CW-CW scenarios. However, simulations considering grain from CW showed a higher climate change impact related to pig production. Compared with the SO-SO scenario, an increase of 17% in climate change impact was found when simulating with soybeans produced in CW, while a 28% increase was observed when simulating with corn and soybeans from CW. As previously stated, the carbon footprint varied among scenarios, mainly due to the impact of transportation and deforestation.

Feeding was the largest source of environmental impact in the pig production scenarios considered in this study, a finding that agrees with several previous studies conducted in Brazil (Spies, 2009; Cherubini *et al.*, 2015a, 2015b; Kebreab, *et al.*, 2016) or other countries (Basset-Mens and van der Werf, 2005; Eriksson *et al.*, 2005). The feed used during the growing-finishing phase was particularly important, accounting for up to 56% of the potential climate change impact, up to 56% of the eutrophication impact, and up to 51% of the acidification impact of finished pigs raised according to the CON feeding program in Brazil. Due to their high contribution to the total environmental impact, feeding practices may be considered a prime target for improvement when developing mitigation strategies for the pig production chain.

### Environmental impact of adopting precision feeding programs

Phase-feeding strategies are the most widely used feeding technique in pig production (Niemi *et al.*, 2010). These feeding programs are designed to maximize animal performance by providing a single feed to all pigs in the herd during a certain period. However, the pigs' nutritional requirements change dynamically over time and also vary greatly among individuals (Pomar *et al.*, 2003; Brossard *et al.*, 2009). By disregarding these variability issues, conventional group phase-feeding programs leads to an inaccurate supply of nutrients, usually with some underfed animals and most pigs receiving more nutrients than they actually need (Hauschild *et al.*, 2010).

The impact of using a mathematical model to estimate real-time daily lysine requirements in a sustainable precision feeding program for growing pigs has been studied in the past. Feeding growing pigs individually with diets adjusted on a daily basis to meet their estimated requirements reduced nutrient intake and excretion without compromising pig performance (Andretta *et al.*, 2014; Andretta *et al.*, 2016). Although the previous results had already indicated some local environmental benefits, the current study pointed out the global impact of adopting precision

feeding practices for growing-finishing pigs.

Applying the PFG program reduced the potential climate change, eutrophication, and acidification impacts compared with the CON program. The greatest mitigation when switching from CON to PFG (equivalent to a 4% savings in potential climate change impact) was observed when pigs were raised in SO on a diet composed of corn cultivated in the same region and soybeans from CW (CW-SO scenario).

Adopting the PFI program further mitigated the environmental impact. Feeding each pig according to its individual nutrient requirements reduced the potential climate change impact by up to 6% and the potential eutrophication and acidification impacts by up to 5% compared with the CON program. Again, the greatest mitigation when switching from CON to PFI was observed in the CW-SO scenario.

It is important to point out that the mitigating impact of precision feeding simulated in this study depends on the nutritional composition of the diet given in the CON treatment. This reference treatment provided all pigs in this group and within each feeding phase with a fixed blend of premixes A and B that had been set based on the requirements of the 80th-percentile pig in the population. This nutritional level had been suggested by previous authors (Hauschild *et al.*, 2010) to maximize the response of the population in terms of body weight gain, and it is in agreement with other results *in vivo* (Brossard *et al.*, 2014) and *in silico* (Brossard *et al.*, 2009).

The proposed precision feeding system represents a paradigm shift in pig production, as it takes into account differences in nutrient requirements among individuals within a population and their dynamics over time. Applying precision feeding techniques significantly improves nutrient-use efficiency (Pomar *et al.*, 2014). Although pig performance (i.e. weight gain) was not changed by the feeding practices under study, some important nutrients are saved by adjusting the provision of nutrients to the dynamic requirements of individual animals. In summary, the environmental benefits of switching from a conventional feeding system to precision feeding programs during the growing-finishing phase came from avoiding nutrient oversupply, changing the ingredient use (feed formulas), and reducing nutrient excretion to the environment.

## Conclusion

This study investigated the global environmental impact of using a mathematical model that estimates real-time daily lysine requirements to deliver a sustainable precision feeding program for growing pigs. Although several studies on the environmental impact of pig production have already been published, only a few of these addressed the effects of changing the feeding program, particularly by using real data (other studies used observations on a population rather than simulated data). In addition, most of these studies were developed for European conditions, and their results cannot be extrapolated to other major pig production regions in developing countries. The results of this study clearly show that precision feeding is an effective approach for improving the environmental sustainability of Brazilian pig farming. In addition, adopting precision feeding techniques for growing-finishing pigs is a highly promising avenue for improving resource-use efficiency (e.g. nutrients) in comparison with conventional group phase-feeding programs.

## References

- AGRINESS (2013). *Melhores da suinocultura - Dados consolidados por estado (6a edição)*. Florianópolis: AGRINESS.
- Alvarenga, R.A.F. (2010). *Avaliação de métodos de AICV: um estudo de caso de quatro cenários de ração para frangos de corte*. Florianópolis: Universidade Federal de Santa Catarina.
- Alvarenga, R.A.F., Silva Júnior, V.P., Soares, S.R. (2012). Comparison of the ecological footprint and a life cycle impact assessment method for a case study on Brazilian broiler feed production. *Journal of Cleaner Production*, 28, 25–32.

- Andretta, I., Pomar, C., Rivest, J., Pomar, J., Lovatto, P.A., Radünz, J. (2014). The impact of feeding growing-finishing pigs with daily tailored diets using precision feeding techniques on animal performance, nutrient utilization, and body and carcass composition. *Journal of Animal Science*, 92, 3925–3936.
- Andretta, I., Pomar, C., Rivest, J., Pomar, J., Radünz, J. (2016). Precision feeding can significantly reduce lysine intake and nitrogen excretion without compromising the performance of growing pigs. *Animal*, 10, 1137–1147.
- Basset-Mens, C., van der Werf, H.M.G. (2005). Scenario-based environmental assessment of farming systems: the case of pig production in France. *Agriculture, Ecosystems & Environment*, 105, 127–144.
- Brossard, L., Dourmad, J.-Y., Rivest, J., van Milgen, J. (2009). Modelling the variation in performance of a population of growing pig as affected by lysine supply and feeding strategy. *Animal*, 3, 1114–1123.
- Brossard, L., Vautier, B., van Milgen, J., Salaun, Y., Quiniou, N. (2014). Comparison of in vivo and in silico growth performance and variability in pigs when applying a feeding strategy designed by simulation to control the variability of slaughter weight. *Animal Production Science*, 54, 1939–1945.
- Centro de Estudos Avançados em Economia Aplicada - Escola Superior de Agricultura "Luiz de Queiroz" (CEPEA-ESALQ) (2014). *Indicadores de preços - CEPEA-ESALQ*, Piracicaba: Universidade de São Paulo.
- Cherubini, E., Zanghelini, G.M., Alvarenga, R.A.F., Franco, D., Soares, S.R. (2015a). Life cycle assessment of swine production in Brazil: a comparison of four manure management systems. *Journal of Cleaner Production*, 87, 68–77.
- Cherubini, E., Zanghelini, G.M., Tavares, J.M.R., Belettini, F., Soares, S.R. (2015b). The finishing stage in swine production: influences of feed composition on carbon footprint. *Environment, Development and Sustainability*, 17, 1313–1328.
- Eriksson, I.S., Elmquist, H., Stern, S., Nybrant, T. (2005). Environmental systems analysis of pig production - The impact of feed choice. *The International Journal of Life Cycle Assessment*, 10, 143–154.
- Ferket, P.R., van Heugten, E., van Kempen, T.A.T.G., Angel, R. (2002). Nutritional strategies to reduce environmental emissions from nonruminants. *Journal of Animal Science*, 80 (E. Suppl. 2), E168–E182.
- Guinée, J.B. (2002). *Handbook on life cycle assessment: Operational guide to the ISO standards*. Amsterdam: University of Amsterdam.
- Hauschild, L., Pomar, C., Lovatto, P.A. (2010). Systematic comparison of the empirical and factorial methods used to estimate the nutrient requirements of growing pigs. *Animal*, 4, 714–723.
- Hauschild, L., Lovatto, P.A., Pomar, J., Pomar, C. (2012). Development of sustainable precision farming systems for swine: Estimating real-time individual amino acid requirements in growing-finishing pigs. *Journal of Animal Science*, 90, 2255–2263.
- Instituto Brasileiro de Geografia e Estatística - IBGE (2014). *Produção Agrícola Municipal de 2012*. Brasília: IBGE.
- Intergovernmental Panel on Climate Change - IPCC (2006). *2006 IPCC guidelines for national greenhouse gas inventories*. Japan: IPCC.
- Kebreab, E., Liedke, A., Caro, D., Deimling, S., Binder, M., Finkbeiner M. (2016). Environmental impact of using specialty feed ingredients in swine and poultry production: A life cycle assessment. *Journal of Animal Science*, 94, 2664–2681.
- Meul M., Ginneberge, C., Van Middelaar, C.E., de Boer, I.J.M., Fremaut, D., Haesaert, G. (2012). Carbon footprint of five pig diets using three land use change accounting methods. *Livestock Science*, 149, 215–223.
- Monteiro, A.N., Garcia-Launay, F., Brossard, L., Wilfart, A., Dourmad, J.-Y. (2016). Effect of feeding strategy on environmental impacts of pig fattening in different contexts of production: evaluation through life cycle assessment. *Journal of Animal Science*, 94, 4832–4847.
- Mosnier, E., van der Werf, H.M.G., Boissy, J., Dourmad, J.-Y. (2011). Evaluation of the environmental implications of the incorporation of feed-use amino acids in the manufacturing of pig and broiler feeds using Life Cycle Assessment. *Animal*, 5, 1972–1983.
- Niemi, J.K., Sevón-Aimonen, M.-L., Pietola, K., Stalder, K.J. (2010). The value of precision feeding technologies for grow-finishing swine. *Livestock Science*, 129, 13–23.
- Pomar, C., Kyriazakis, I., Emmans, G.C., Knap, P.W. (2003). Modeling stochasticity: Dealing with populations rather than individual pigs. *Journal of Animal Science*, 81, E178–E186.
- Pomar, C., Hauschild, L., Zhang, G.-H., Pomar, J., Lovatto, P.A. (2009). Applying precision feeding techniques in growing-finishing pig operations. *Revista Brasileira de Zootecnia*, 38, 226–237.
- Pomar, C., Pomar, J., Rivest, J., Cloutier, L., Letourneau-Montminy, M.-P., Andretta, I., Hauschild, L. (2014). Estimating real-time individual amino acid requirements in growing-finishing pigs: Towards a new definition of nutrient requirements in growing-finishing pigs? In N.K. Sakomura, R.M. Gous, I. Kyriazakis, L. Hauschild (Eds.), *Nutritional modelling in pigs and poultry* (pp. 157–174). Wallingford: CABI.
- Prudêncio da Silva, V., van der Werf, H.M.G., Spies, A., Soares, S.R. (2010). Variability in environmental impacts of Brazilian soybean according to crop production and transport scenarios. *Journal of Environmental Management*, 91, 1831–1839.
- Rostagno, H.S., Albino, L.F.T., Donzale, J.L., Gomes, P.C., Oliveira, R.F., Lopes, D.C., Ferreira, A.S., Barreto, S.L.T., Euclides, R.F. (2011). *Tabelas Brasileiras para Aves e Suínos*. Viçosa: Universidade Federal de Viçosa.
- Spies, A. (2009). Avaliação de impactos ambientais da suinocultura através da análise de ciclo de vida - ACV. In C.R. Miranda, M. Miele (eds.), *Suinocultura e meio ambiente em Santa Catarina: indicadores de desempenho e avaliação*. **Proceedings of the 14<sup>th</sup> International Conference on Precision Agriculture** June 24 – June 2, 2018, Montreal, Quebec, Canada

- sócio-econômica (pp. 13–43). Concórdia: Embrapa Suínos e Aves.
- Talamini, D.J.D., Martins, F.M., Arboit, C., Wolozsim, N. (2006). Custos agregados da produção integrada de suínos nas fases de leitões e de terminação. *Custos e Agronegócio*, 2, 64–83.
- van der Werf, H.M.G., Petit, J., Sanders, J. (2005). The environmental impacts of the production of concentrated feed: the case of pig feed in Bretagne. *Agricultural Systems*, 83, 153–177.
- van Zeist, W.J., Marinussen, M., Broekema, R., Groen, E., Kool, A., Dolman, M., Blonk H. (2012a). *LCI data for the calculation tool Feedprint for greenhouse gas emissions of feed production and utilization: Crushing industry*. Gouda: Blonk Consultants.
- van Zeist, W.J., Marinussen, M., Broekema, R., Groen, E., Kool A., Dolman, M., Blonk, H. (2012b). *LCI data for the calculation tool Feedprint for greenhouse gas emissions of feed production and utilization: Dry milling industry*. Gouda: Blonk Consultants.
- Wathes, C.M., Kristensen, H.H., Aerts, J.M., Berckmans, D. (2008). Is precision livestock farming an engineer's daydream or nightmare, an animal's friend or foe, and a farmer's panacea or pitfall? *Computers and Electronics in Agriculture*, 64, 2–10.